



Strasbourg, 9 June 2010  
[Inf09e\_2010]

**T-PVS/Inf (2010) 9**

CONVENTION ON THE CONSERVATION OF EUROPEAN WILDLIFE  
AND NATURAL HABITATS

**Standing Committee**

30<sup>th</sup> meeting  
Strasbourg, 6-10 December 2010

---

## **CLIMATE CHANGE AND THE BIODIVERSITY OF EUROPEAN ISLANDS**

**- First Draft -**  
**8<sup>th</sup> June 2010**

*Document prepared by  
Ms Cordula Epple, UNEP-WCMC*

---

*This document will not be distributed at the meeting. Please bring this copy.  
Ce document ne sera plus distribué en réunion. Prière de vous munir de cet exemplaire.*

**TABLE OF CONTENTS**

<b>Executive summary</b> .....	3
<b>1. Introduction</b> .....	3
<b>2. Island biodiversity in Europe</b> .....	4
a. General features of island biodiversity.....	4
b. Geographic overview of European islands and their main characteristics .....	4
c. Importance of islands for biodiversity in Europe .....	9
d. State of conservation efforts on European islands .....	10
<b>3. Impacts of climate change on the biodiversity of European islands</b> .....	10
a. Observed impacts .....	10
b. Projected impacts.....	11
<b>4. Implications of climate change for conservation efforts on European islands and options for action</b> .....	15
<b>5. Research needs</b> .....	16
<b>6. Conclusion and recommendations</b> .....	16
<b>7. Acknowledgements</b> .....	17
<b>8. References</b> .....	17
<b>Appendices</b> .....	19

## EXECUTIVE SUMMARY

European islands are home to many species and habitats of conservation concern, including endemic as well as threatened biodiversity. Endemism is largely concentrated on islands in the Mediterranean and Macaronesian region. There are significant knowledge gaps concerning current and potential future impacts of climate change on European island biodiversity. However, there is enough evidence to demonstrate that impacts already take place and are likely to increase in future. Processes related to climate change which are particularly relevant in the island context include sea level rise and the possibility of increasing incidence of invasive alien species. Available measures to support adaptation for biodiversity are similar to those recommended for other areas. However, possibilities to enhance connectivity beyond the individual island are limited. Ex situ and translocation measures might be considered for endemic island taxa where no other options exist. Priority attention should be given to islands in the Mediterranean and Macaronesian region both because of their high endemism and because of expected changes in precipitation patterns, and within these islands to endemic species which are already considered threatened. Further efforts in monitoring and research are recommended.

### 1. INTRODUCTION

The specific environmental and socio-economic conditions of islands provide both challenges and opportunities for nature conservation. A targeted approach to biodiversity issues which takes these characteristic features into account has recently been promoted through the adoption of the Programme of Work on Island Biodiversity under the Convention on Biological Diversity (CBD) in 2006, as well as the formation of the Global Islands Partnership (GLISPA)<sup>1</sup>.

In the CBD Programme of Work, climate change is pointed out as one of the important threats to island biodiversity and all Parties are encouraged to implement adaptation measures and strategies to strengthen the resilience of species and ecosystems, and to carry out research and monitoring to improve the knowledge base for such activities.

From another perspective, the available literature on climate change impacts on ecosystems frequently mentions islands as an area of high concern (e.g. Berry 2008). Reasons for this include the high ratio of coastline to overall land area and the ensuing threat of habitat loss through sea level rise as well as the fact that for many island populations of plant and animal species it will be even more difficult than for mainland populations to adapt to climate change by dispersing into newly suitable areas.

Because of the wealth of islands that is present on the European continent, these findings should be reflected in any regional efforts to conserve biodiversity in the face of rapid global change.

The present study was prepared on behalf of the Council of Europe for consideration by the Group of Experts on Biodiversity and Climate Change which has been established under the Bern Convention in 2006. It aims to outline the state of knowledge concerning island biodiversity in Europe and the likely impacts of climate change on its components, and to provide a starting point for discussion on actions to be taken by Parties. It builds upon previous reports reviewed by this Group of Experts which have dealt with other aspects of the link between climate change and the conservation of European species and habitats<sup>2</sup> as well as the guidance provided to Parties on the issue of biodiversity and climate change in Recommendations No. 135 (2008) and No. 143 (2009), and is intended to complement the ongoing work of the Groups of Experts on European Island Biological Diversity and Invasive Alien Species.

In line with the agreement made by the Group of Experts on Island Biological Diversity, the scope of this report has been set to include the Macaronesian islands, but not the overseas territories of European

---

<sup>1</sup> see <http://www.cbd.int/island/glispa.shtml>

<sup>2</sup> Huntley (2007, Doc. T-PVS/Inf (2007) 3), Ferrer et al. (2008, Doc. T-PVS/Inf (2008) 1 rev), Capdevila-Arguelles & Zilletti (2008, Doc. T-PVS/Inf (2008) 5 rev.), Berry (2008, Doc. T-PVS/Inf (2008) 6 rev.), Henle et al. (2008, Doc. T-PVS/Inf (2008) 11 rev.), Harley (2008, Doc. T-PVS/Inf (2008) 12 rev.), Wilson (2009, Doc. T-PVS/Inf (2009) 8), Heywood & Culham (2009, Doc. T-PVS/Inf (2009) 9), Araujo & Garcia (2009, Doc. T-PVS/Inf (2009) 10)

countries in the Pacific and Indian Oceans or the Caribbean Sea. Only marine islands have been considered.

## **2. ISLAND BIODIVERSITY IN EUROPE**

### **a. General features of island biodiversity**

The characteristics of the species assemblages found on islands are determined by many factors, including size, age, distance to other islands and the mainland, climatic history, current climate, relief and geology. Thus, although as a general rule islands are poorer in species than comparable mainland areas (Whittaker and Palacios 2007), their biodiversity often exhibits unique features and a high degree of endemism, from the genetic to the ecosystem level.

A useful distinction can be made between remote islands for which evolution has in the past been a faster process than immigration – termed “oceanic” islands by Whittaker and Palacios (2007, with reference to Williamson 1981) because many areas which fit this description are located off the continental shelf – and islands in relative proximity to the mainland, which are characterised by higher immigration rates and may be called “continental”. Since both the evolution and immigration rates vary between different groups of organisms, there is a continuous range of intermediate forms between these two types. For example, many insular locations may have bird populations which are closely connected to those of the nearby continent, while their terrestrial mammal populations are effectively isolated.

The significance of islands for global biodiversity conservation has been highlighted by Fonseca et al. (2006). Although islands make up only some 5 % of the global land area, their endemic biota are estimated to include about 20 % of the world’s vascular plant species and 15 % of all mammal, bird and amphibian species. Endemism is often correlated with vulnerability to various factors of threat, and island endemites account for around one third of the mammal, bird and amphibian species which have been classified as threatened on a global scale.

Endemism is, however, not the only factor which contributes to the high conservation value of many islands. The array of threats or pressures acting on a species in a given location is likely to differ from the mainland, and elements such as land use intensity or predator pressure may be either more or less pronounced in each individual island’s setting. Thus, vulnerable species which have been lost from much of their former distribution area may find refugial niches and persist in significant numbers on islands with suitable circumstances. Ground-breeding bird and reptile species often form large colonies on inaccessible or predator-free island sites. Islands located along migration pathways may also provide important stopover and resting sites for terrestrial animals.

Threatening factors which often have more severe effects in island ecosystems both because of the existing space limitations and the specialised nature of island biota include habitat destruction, uncontrolled tourism development, invasive alien species, and climate change (Orueta 2009, Carnevali & Genovesi 2009).

In line with the above-mentioned facts, many of the lists of priority sites for conservation which have been developed by different international organisations (e.g. the Global 200 Ecoregions (WWF), the Endemic Bird Areas (BirdLife International), the Biodiversity Hotspots (Conservation International) and the AZE sites (Alliance for Zero Extinction)) include a high proportion of sites situated entirely on islands (one quarter to about half of the sites for the listings cited above) as well as numerous sites comprising mainland areas as well as islands (Fonseca et al. 2006).

### **b. Geographic overview of European islands and their main characteristics**

Islands are a prominent feature of the European continent with its long and irregular coastline. Several European countries are entirely situated on islands, and except for a small number of landlocked states almost all countries have islands within their territories. As stated by Carnevali and Genovesi (2009), although an inventory of European islands has not been completed yet, the Global Island Database

(<http://gid.unep-wcmc.org/>)<sup>3</sup> holds information about more than 50,000 islands belonging to Europe. Of these, around 500 are larger than 20 km<sup>2</sup>, and their total area is more than 700,000 km<sup>2</sup>, or more than 7 % of the surface of Europe (Orueta 2009).

There is an enormous diversity among the European islands as regards their size, geology, morphology and climate, such that any kind of general statement about their characteristics would be impossible. Typical 'island phenomena' such as a strong maritime influence, resource limitations and a high share of specialized biota and communities are more prevalent on smaller and more isolated territories, while the bigger islands such as Great Britain have more commonalities with coastal states on the mainland. Table 1 gives an overview of major islands and archipelagos in the biogeographical regions of Europe and some of their characteristic features, referring to the delineation of biogeographical regions as defined by the European Commission and the Council of Europe for the purpose of evaluation and reporting on nature conservation (EEA 2002-2008).

---

<sup>3</sup> The Global Island Database was initiated within the framework of the Global Island Partnership and is being developed in a multilateral partnership project led by UNEP-WCMC.

**Table 1: Major islands and archipelagos in the biogeographical regions of Europe and some of their characteristic features (Sources: Global Island Database<sup>4</sup>, UNEP Island Directory<sup>5</sup>, EEA 2002-2008, Wikipedia)**

Biogeographic region	Name of island or archipelago <sup>6</sup>	Country or countries <sup>7</sup>	Characteristics
Arctic	Svalbard	Norway	Largest island Spitsbergen; total land area of archipelago 61,022 km <sup>2</sup> ; distance to mainland (Norway) 660 km; highest elevation 1,713 m asl
	Iceland	Iceland	Land area 101,794 km <sup>2</sup> ; distance to mainland (Norway) 970 km; main vegetation type tundra; highest elevation 2,110 m asl; coasts consisting mainly of rock and sand with open fjords
	Coastal archipelago of Northern Norway	Norway	Largest island Hinnoya, with an area of 2,204 km <sup>2</sup> ; large number of islands of different sizes at short distance from mainland
Boreal	Aland islands and islands of the Finnish Archipelago Sea	Finland	Largest island Fasta Aland, with an area of 879 km <sup>2</sup> ; large number of islands of different sizes at short distance from mainland, total area about 2,000 km <sup>2</sup> ; more than 90 % of the territory less than 30 m asl; geology characterised by bedrock
	Hiiumaa	Estonia	Land area 1,026 km <sup>2</sup> ; highest elevation 68 m; distance to mainland 22 km; includes calcareous soils
	Saaremaa	Estonia	Land area 2,680 km <sup>2</sup> ; highest elevation 54 m; distance to mainland 8 km; includes calcareous soils
	Gotland	Sweden	Land area 3,015 km <sup>2</sup> ; highest elevation 82 m; distance to mainland 90 km; includes calcareous soils
	Swedish coastal archipelago on the Baltic Sea <sup>8</sup>	Sweden	Largest island group Stockholm archipelago, with an area of about 530 km <sup>2</sup> ; large number of islands of different sizes at short distance from mainland;

<sup>4</sup> <http://gid.unep-wcmc.org>

<sup>5</sup> <http://islands.unep.ch/>

<sup>6</sup> Note that the designations used in this report are not meant to express any opinion concerning the legal status of any country or island territory.

<sup>7</sup> Several island territories have some form of political autonomy within their countries, thus the information in this column should not be taken as a description of political status

		more than 90 % of the territory less than 30 m asl; geology characterised by bedrock
	Swedish and Norwegian coastal archipelagos on the Skagerrak Sea	Sweden / Norway
	Norwegian archipelago on the western coast south of the Polar Circle	Norway
<b>Atlantic</b>	Faroe Islands	Denmark
	British Isles and Ireland	United Kingdom, Ireland
	Frisian Islands	Denmark, Germany, Netherlands
<b>Continental</b>	Danish Islands in the Kattegat and Baltic Sea (incl. Zealand, Fyn, Bornholm)	Denmark
	Swedish coastal archipelago in the Kattegat and Southern Baltic Sea up to Oland	Sweden
	Oland	Sweden
	Islands on the Pomeranian Bay (incl. Rugen, Usedom, Wolin)	Germany, Poland
<b>Mediterranean</b>	Balearic Islands	Spain

<sup>8</sup> with the exception of the small area south of Oland which belongs to the Continental region

Corsica	France	Land area 8,741 km <sup>2</sup> ; highest elevation 2,706 m; distance to mainland 90 km
Sardinia	Italy	Land area 23,962 km <sup>2</sup> ; highest elevation 1,834 m; distance to mainland 200 km
Sicily	Italy	Land area 25,531 km <sup>2</sup> ; highest elevation 3,320 m; distance to mainland 3 km
Malta	Malta	Total land area of archipelago 316 km <sup>2</sup> ; highest elevation 253 m; distance to mainland 220 km
Adriatic islands	Croatia, Montenegro, Albania, Italy	Largest island Cres with a total land area of 406 km <sup>2</sup> ; highest elevation 778 m; large number of islands at short distance from mainland
Ionian islands	Greece	Total land area 2,307 km <sup>2</sup> ; highest elevation 1,630 m; distance to mainland less than 1 km
Aegean islands (incl. Crete)	Greece, Turkey	Largest island Crete with a total land area of 8,336 km <sup>2</sup> ; highest elevation 1,611 m; large number of islands at varying distance to mainland
Cyprus	Cyprus	Land area 9,317 km <sup>2</sup> ; highest elevation 1,952 m; distance to mainland 75 km
<b>Macaronesian</b> Azores	Portugal	Total land area 2,346 km <sup>2</sup> ; highest elevation 2,351 m; distance to mainland 1,390 km; volcanic origin
Madeira	Portugal	Total land area 828 km <sup>2</sup> ; highest elevation 1,826 m; distance to mainland 520 km; volcanic origin
Canary Islands	Spain	Total land area 7,447 km <sup>2</sup> ; highest elevation 3,718 m; distance to mainland 97 km; volcanic origin



### c. Importance of islands for biodiversity in Europe

The conservation assets of European islands are as diverse as their physiogeographic and socioeconomic characteristics. Many islands especially in those regions which are easily accessible and attractive for tourism or leisure activities are suffering from overexploitation of scarce resources and destruction of coastal habitats, whereas others still hold a large extent of relatively unimpacted ecosystems.

Because of their high ratio of coastline to interior, islands are hosting a significant share of the total area of coastal, littoral and shallow water marine habitats in Europe. For example, although islands make up only 5.8 % of the territory of Croatia, they account for about 70 % of the Croatian coastline (source: Croatian national report to the Bern Convention on activities related to Biological Diversity on European islands<sup>9</sup>). For the whole of the Mediterranean Sea, according to EEA (2002-2008), island coasts make up about 40 % of the total coastline. Coastal and halophytic communities represent a large share of the Bern Convention habitats as listed in Resolution 4 (96). The machair, a specific type of coastal grassland with a long history of extensive agricultural use, is only found on the Hebrides as well as the northwest coasts of Ireland and Scotland. Although marine habitats such as seagrass meadows or mussel beds around islands are not generally distinct from those of the mainland coast, they may be better preserved in many areas due to a lower degree of disturbance.

As stated above, one of the main reasons for paying particular attention to islands in biodiversity conservation is their high level of endemism. Although many endemic species have been lost from European islands as a consequence of human settlement (Orueta 2009), the number of species confined to one or several islands is still significant. For example, according to Delanoe et al. (1996), the rate of endemic species among vascular plants on the larger Mediterranean islands is on average around 10 %. Even higher levels of endemism are reached in the Macaronesian islands with their unique blend of North Atlantic, African and Mediterranean biogeographic influences and special vegetation types such as the humid evergreen laurel forest and the dry Canary pine forests.

There is currently no complete inventory of endemic species on European islands (Orueta 2009), and new taxa are still being discovered. However, some of the available information on well-researched taxonomic groups has been compiled in annexes 1 and 2. These tables clearly demonstrate the concentration of endemic species in the Mediterranean and Macaronesian regions. The near complete absence of species-level endemism from northern European islands is believed to be largely a consequence of their recent glaciation history (EEA 2002-2008).

Another prominent feature from a conservation point of view is the importance of many islands for breeding and migrating birds as well as marine mammals and reptiles. Both the United Kingdom and Ireland hold a large number of internationally important wintering, stopover and breeding sites for birds of conservation concern on the main islands as well as the numerous smaller islands. 31 of the Special Protection Areas classified under the EU Birds Directive in the United Kingdom are situated on offshore islands, accounting for some 3,788,000 breeding seabirds<sup>10</sup>. As noted in the French national report on island biodiversity, islands are hosting the main part of the national population of several seabird species, e.g. Cory's Shearwater (*Calonectris diomedea*), Storm petrel (*Hydrobates pelagicus*) and the subspecies of the European Shag *Phalacrocorax aristotelis subspp. desmaretii*.

The importance of Arctic islands as one of the world's most important reproduction areas for seabirds is pointed out in EEA (2002-2008), highlighting also the role of seabird populations in the region's ecosystem. In the same publication, the role of the archipelagos along the Swedish and Finnish Baltic coasts as breeding sites for large numbers of ducks and waders is also emphasized. For example, the small islands and skerries are reported to contain an estimated 600,000 pairs of eider ducks. Also, all three species of seal found in the Baltic - the grey seal (*Halichoerus grypus*), the harbour (common) seal (*Phoca vitulina*) and the Baltic ringed seal (*Phoca hispida botnica*) - live mainly in the archipelagos. Migratory waterbirds are increasingly wintering in the lagoons and shallow waters of the Southern Baltic, and the extensive mudflats and shallow waters of the Wadden Sea area are used as stopover or wintering sites by more than 12 million birds per year, while also

<sup>9</sup> Doc. T-PVS/Inf(2009)12

<sup>10</sup> Doc. T-PVS/Inf(2009)12

hosting marine mammals such as the harbour seal (*Phoca vitulina*), the grey seal (*Halichoerus grypus*) and the bottle-nose dolphin (*Tursiops truncatus*) (EEA 2002-2008).

In addition to coastal communities, several other Bern Convention habitats, e.g. European wet heaths and Mediterranean shrub formations, are commonly found on islands, and a few are endemic to specific islands or island groups (e.g. Macaronesian heaths and certain types of Aleppo pine woods). Orueta (2009) has proposed a more thorough assessment of the distribution of habitats on European islands and their conservation status as a potential line of work for the Bern Convention Group of Experts on Island Biological Diversity.

#### **d. State of conservation efforts on European islands**

Because of the importance of Mediterranean islands as a biodiversity hotspot, several international research and conservation efforts have focussed in this region, for example the work of the Mediterranean Island Plant Specialist Group of IUCN<sup>11</sup>.

Although the work of the Global Island Partnership has in the past focussed mainly on tropical areas, there are some activities in the European region as well. For example, Croatia has made a commitment to work towards an improvement of the network of Marine Protected Areas in the Mediterranean by 2012.

On the national level, the designation of protected areas, notably including the sites of the Natura 2000 network, has been one of the focal areas of conservation work in the past decades. In many cases, the share of protected areas is higher on the island part of a country's territory than on the mainland. For example, more than 85 % of the Croatian island area and 40 % of the area of the Canary Islands has been included in the Natura 2000 network (national reports to the Bern Convention on activities related to Biological Diversity on European islands<sup>12</sup>). More than half of the area of the Svalbard archipelago is included in protected areas under Norwegian law (EEA 2002-2008). According to Orueta (2009), about 10 % of the European Biosphere Reserves are situated on islands. However, island protected areas often face particular logistic constraints as compared to the mainland.

### **3. IMPACTS OF CLIMATE CHANGE ON THE BIODIVERSITY OF EUROPEAN ISLANDS**

#### **a. Observed impacts**

Observations of climate change impacts on the biodiversity of European islands are anecdotal in nature, and no attempt could be made to locate all items of literature. Most of the readily available information relates to well-studied groups of species and larger, densely populated islands with a tradition of species monitoring, while knowledge on species occurrence and trends is generally poor on smaller islands due to logistical barriers. Based on a comprehensive review of evidence for the impacts of climate change on biodiversity in the European Union, Hodgson et al. (2009) concluded that there is a marked imbalance of information coverage in all of the European biogeographical regions.

The types of changes which have been most frequently recorded and analysed include phenological changes and expansions in distribution of conspicuous species such as birds or butterflies. However, from a practical conservation point of view, information on overall population trends and changes in distribution areas especially for rare or declining species is more urgently needed.

Hodgson et al. (2009) cite a large number of studies demonstrating shifts in phenology and distribution ranges of birds, butterflies and other terrestrial and littoral invertebrates in Great Britain, while stating that for other parts of the Atlantic biogeographical region, literature is more scarce.

Monitoring data on moths in Iceland (time series beginning in 1995) show a trend towards earlier spring adult appearance (Nordic Council of Ministers 2009).

---

<sup>11</sup>

[http://www.iucn.org/about/work/programmes/species/about\\_ssc/specialist\\_groups/directory\\_specialist\\_groups/directory\\_sg\\_plants/ssc\\_mediterranean\\_island\\_plant/](http://www.iucn.org/about/work/programmes/species/about_ssc/specialist_groups/directory_specialist_groups/directory_sg_plants/ssc_mediterranean_island_plant/)

<sup>12</sup> Doc. T-PVS/Inf(2009)12 and T-PVS (2009) 13

Observations of the reproductive success, survival rates and population numbers of seabirds breeding on islands around the North Sea and the North Atlantic Ocean have shown large-scale decreases in recent decades. These are attributed to a combination of factors, notably overfishing and climate change both leading to a decline in food sources. For example, a decrease in the population of the thick-billed murre (*Uria lomvia*) and fulmar (*Fulmarus glacialis*) has been observed in Iceland for the period from the 1980s to 2005 (Nordic Council of Ministers 2009).

Changes in the behaviour of migratory birds have been recorded for several species since the last century. These include not only changes in timing, but also changes to migratory routes and destinations, which means that the relative importance of islands for the survival of certain species may increase or decline. Ferrer et al. (2008) have reviewed the evidence for climate change effects on migration routes of European birds. One of the most well known examples of recent changes in migratory direction is that of the Blackcap (*Sylvia atricapilla*), whose winter distribution has now largely shifted from southern Europe to the British Islands.

As an example of an island-based study focussing on impacts on vegetation, del Arco (2006, as cited in Hodgson et al. 2009) found evidence for decreased plant productivity linked to reduced rainfall in the Canary islands.

Due to the obvious bias in literature coverage, no comparisons can be made about the degree of already observed climate change impacts on the biodiversity of different island regions.

## **b. Projected impacts**

One of the challenges with regard to determining potential climate change impacts on islands lies in the fact that the coarse spatial resolution of many currently available projections of future changes in climatic parameters makes it difficult to derive valid statements for many of the smaller islands. The need for downscaled projections to support assessments of potential climatic changes on small islands has also been noted by Christensen et al. (2007) in the IPCC 4<sup>th</sup> Assessment Report.

In spite of these limitations, some basic assumptions can be drawn from available data on projected climatic developments at the European scale as well as from regional models that have been developed for certain areas. Generally, increases in temperature are expected to be attenuated on islands as compared to continental areas due to the buffering effect of the ocean. In line with this, according to an analysis of the temperature changes which might affect Natura 2000 sites in Europe (Bertzky et al. 2009, based on data derived from the HadCM3 model), even under the higher A2 scenario most Natura 2000 sites situated on islands in the Mediterranean, Atlantic and Continental biogeographical regions would only experience a warming of 1-2° C during the second half of the century, while the largest share of the mainland part of the European Union would see temperatures rise by 2-3° C. In the northwestern part of the British Isles and Ireland, the increase in mean annual temperature is even projected to remain below 1° C. However, it has to be noted that the scale of the oceanic effect might have been overestimated in this study due to a low data resolution, as the climatic data retrieved from the IPCC Data Distribution Centre was used without further regionalisation.

With regard to precipitation, the data presented by EEA (2007, with reference to Schroter et al. 2005) suggests more variation among the biogeographical regions. The projections for changes in annual precipitation during the 21<sup>st</sup> century (again for the A2 scenario and based on the HadCM3 model) show an increase of 10-15 % for all parts of the Norwegian archipelago (both within the Arctic as well as the Atlantic and Continental regions), as well as for most islands within the Boreal and Continental regions. In the Atlantic region, only for Ireland and Great Britain more differentiated shifts in rainfall are projected, ranging from an increase of 0-5% in the Southeast of Great Britain to increases of up to 15-20% on the eastern coast of Scotland. Only some of the islands in the Mediterranean region would experience an overall decrease in annual rainfall over parts of their area, with reductions of up to 10-15% for Sardinia and Sicily. Because of the shift of precipitation towards the winter months which is expected for all of Europe (Christensen et al. 2007), these average values imply a significantly drier summer climate in the Mediterranean region. Consequences would include not only prolonged periods of low soil moisture which would affect vegetation, but also an increasing risk of wildfires, which are already an issue of serious concern in many Mediterranean islands. As

stated in the IPCC 4<sup>th</sup> Assessment Report, the Mediterranean region as a whole is likely to be seriously impacted by climate change as a consequence of increasing drought and heat stress.

More detailed regional projections of climatic changes are available mostly for larger island countries rather than single islands within a country's territory.

In order to gain further insights into the likely impacts of future climate change on species and habitats, various approaches can be used. In recent years, a number of studies have used climate envelope models in order to analyze how the potentially suitable climate space of species from different taxonomic groups might shift geographically as a consequence of climate change (e.g. Berry et al. 2005; Berry et al. 2007; Araujo et al. 2006; Thuiller et al. 2005; Huntley et al. 2007, Settele et al. 2008).

However, these approaches can only be applied to species for which a minimum amount of distribution and ecological data is available. This is often not the case for rare species and species of less well researched taxonomic groups. Another important limitation with regard to application in an island context lies in the fact that climate envelope modelling is based on the assumption that species are currently filling the whole of their suitable climatic niche. For narrowly endemic species, whose distribution is likely to be largely determined by historical factors and dispersal barriers (both to the species itself and to potential predators and competitors), this might justify special caution when interpreting the results<sup>13</sup>.

Finally, the problem of scale which was referred to above with regard to available climate projections also applies to climate envelope modelling. Even where finer scale data are available for projections of climate parameters, their application in simulations of the potential future ranges of species may be limited by the resolution of available data on current species distributions (cp. Thuiller et al. 2005, Huntley et al. 2008). Climatic niches which are only realized on small areas would not be captured by the models.

In order to assess the vulnerability<sup>14</sup> of island taxa to climatic change, other approaches including expert opinion based on biological and ecological information and a structured analysis of species' life history and ecological traits should therefore be considered. Work on identifying biological traits which could indicate high susceptibility to climate change and a trait-based assessment of species vulnerabilities to climate change for a large number of bird, amphibian and reef-building warm water corals have been carried out in a study led by IUCN (Foden et al. 2008). The final outcomes of this assessment are yet to be published, but could provide guidance for future work.

Sajwaj et al. (2009) have combined data derived from climate envelope modelling with information on species characteristics that can be used to gauge adaptive capacity in order to develop a framework for assessing the vulnerability of species which are protected under the Habitats and Birds Directives of the European Union. Vulnerability scores could be produced for a total of 212 (~ 24.4%) out of 869 species listed in the Birds and Habitats Directives. Of these, 149 were bird species, 12 amphibian species, 12 reptile species, 13 butterfly species and 26 plant species. For the remaining 657 (869 minus 212) species of Community Interest no suitable modelling data was available.

135 species were assessed in the study as showing a high, very high, critical or extremely critical vulnerability under at least one of the examined climate scenarios and time horizons, leaving only 77 species with moderate to low vulnerability. The authors attribute this to the fact that most species protected under the EU Nature Directives are already threatened or rare and have specific habitat requirements. They point out that constraints to species' ability to move to and colonise new areas with suitable climate (e.g. because of limited dispersal abilities, lack of suitable habitat, or low levels of emigration due to small population sizes etc) are the main factor contributing to vulnerability.

---

<sup>13</sup> For a more detailed discussion of the methodological limitations of climate envelope modelling, see also Heywood & Culham (2009).

<sup>14</sup> The term vulnerability is used here (in line with the definition provided in the IPCC 4<sup>th</sup> Assessment Report) to describe the degree to which a species is likely to be adversely impacted by climate change when its de facto exposure to changes in climatic parameters, its sensitivity and its adaptive capacity are all taken into account.

Although very few of the species assessed in the study are endemic to islands (which may be explained by the lack of data for use in modelling), a significant number also occur on islands and could be given special attention in efforts to protect island biodiversity against negative impacts of climate change. Annex 3 to this document contains a list of Bern Convention species which have been identified by Sajwaj et al. as being at least highly vulnerable.

In contrast to the large body of literature concerning climate change impacts on individual species or species groups, relatively few authors have attempted to address the potential impacts on the level of habitats and ecosystems. Since models for the projection of ecological interactions between species (such as competition between different plant functional types) are highly complex and currently only applicable to very limited sets of species or species groups (Kuhn et al. 2009), it is hardly possible to model the future species assemblages that would determine the characteristics of the ecosystems present in a certain location. The vulnerability of habitats and ecosystems has therefore often been considered through alternative approaches, including expert judgment and the use of information on the vulnerability of characteristic plant and animal species as a surrogate (Sajwaj et al. 2009).

So far no comprehensive analyses on the vulnerability of Bern Convention habitats to climate change have been carried out, and the amount of research and consultations that would be necessary for such a study is far beyond the scope of this paper. Therefore only some specific considerations which appear relevant to the context of European islands can be presented here.

As can be seen from table 5, the largest number of vulnerable species as assessed by Sajwaj et al. (2009) occur on islands in the Mediterranean region. There are several possible reasons for this, including a correlative effect of the higher species richness of the Mediterranean region as compared to other parts of Europe, and the projected severe impacts of climate change in the Mediterranean region as a whole. It should thus not be readily interpreted as an indication of a particularly high vulnerability of Mediterranean island ecosystems.

On the other hand, the high degree of endemism on the islands of the Mediterranean and Macaronesian regions does seem to imply a special concern.

Table 5: Current occurrence of Bern Convention species assessed by Sajwaj et al. (2009) as being at least highly vulnerable to climate change on islands in the different biogeographical regions of Europe (as assessed by the underlying atlas data)

<b>Biogeographical region</b>	<b>Number of species</b>
Arctic	15
Atlantic	34
Boreal	30
Continental	20
Mediterranean	58
Macaronesian	2

An important factor to be kept in mind when thinking about the prioritization of conservation efforts is the likelihood that as species distributions shift with climate change, the relative importance of certain island regions for biodiversity conservation may change. For example, if the projected shifts in suitable climate space as presented in the Climatic Atlas of European Breeding Birds (Huntley et al. 2008) are considered for the vulnerable species listed in Annex III, the number of species which could potentially occur on islands in the Arctic region would double from 14 to 28 by the end of this century.

One specific factor associated with climate change that will have a particularly severe effect on islands is the expected continuation of global sea level rise. In many areas, this will add to the already high pressure on coastal and littoral habitats due to settlement and infrastructure construction. Only for some parts of Scandinavia, sea level rise is expected to be counteracted by the ongoing process of postglacial land uplift (HELCOM 2007).

As noted by Orueta (2009), impacts will be caused not only by changes in mean sea level but also by changes in the frequency of extreme events such as storm surges. According to the IPCC 4<sup>th</sup> Assessment Report, several studies have projected climate-driven increases in extreme water levels in

the European shelf region, although Christensen et al. (2007) state that estimates of future coastal flooding still hold substantial uncertainty. Soft substrate islands, e.g. the islands lining the coasts of the Wadden Sea and the southern Baltic Sea which are largely made up of glacial sediments, are facing a particular risk of high losses in land area due to increased coastal erosion.

Berry et al. (2007) have assessed the potential impacts of sea level rise on intertidal coastal habitats in Europe, concluding that the most vulnerable intertidal habitats are those around the Black, Mediterranean and Baltic Seas. This is explained by the fact that these seas have low tidal ranges and thus coastal ecosystems are more sensitive to sea level rise.

Climate change is also expected to increase the likelihood of threats to native island biota caused by invasive alien species<sup>15</sup>. Already now, invasive species are having particularly detrimental effects on island biodiversity in Europe as well as other regions of the world, despite significant efforts at control and eradication (Genovesi & Shine 2003; Carnevali & Genovesi 2009).

Examples include wild boar (*Sus scrofa*) introduced as game animals on some Croatian islands, rats (*Rattus rattus*) and exogenous fish parasites on Corsica, *Opuntia ficus-indica* on the Italian Ponziene islands, mink (*Mustela vison*) and raccoon dog (*Nyctereutes procyonoides*) on Swedish Baltic Sea islands and the macroalga *Gracilaria vermiculophylla* in the intertidal zone of the Gothenburg archipelago, and *Rhododendron ponticum* in Ireland (Source: national reports to the Bern Convention on activities related to Biological Diversity on European islands<sup>16</sup>).

Many island communities are susceptible to invasion because they have developed under conditions of reduced competition and predation. Climate change is expected to trigger a significant rise in the number of alien species becoming invasive as well as to favour the spread of many invasive alien species that are already established (Capdevila-Arguelles & Zillett 2008). Invasive species may be more able than native ones to benefit from the changing environmental conditions including higher temperatures, decreasing numbers of frost days and elevated levels of carbon dioxide. However, predictions about invasion success at species level are difficult to make.

More frequent extreme events such as storms, floods, droughts or fire may also increase the susceptibility of ecosystems to invasion. Gritti et al. (2006, as cited by Heywood & Culham 2009) have studied the vulnerability of ecosystems in five Mediterranean islands (Mallorca, Corsica, Sardinia, Crete and Lesvos) to climate change and invasion by exotic plants. Their simulations predict that while climate change on its own would not have any severe impacts, as an effect of habitat disturbance in the longer term almost all the ecosystems would be dominated by invasive aliens.

Some work on identifying priority areas for efforts to control the impacts of invasive alien species on European islands, including the development of a European Invasive Alien Species Island Information System, has been initiated under the Bern Convention (Carnevali & Genovesi 2009)<sup>17</sup>.

Finally, climate change may also lead to significant indirect impacts on biodiversity, for example due to adaptations in land use or to mitigation activities such as the increased use of renewable energies. With regard to the latter, the development of wind farms has in recent decades been an issue of concern for conservationists in many island regions, and further expansion is to be expected.

Island-specific projections or expert opinions on impacts are available only for certain regions and taxa, again with a heavy bias towards the British Isles. One of the earliest and most comprehensive studies examining potential climate change impacts was the MONARCH project, which focussed on Britain and Ireland. Among other results, montane heaths, native pine woodland and peat bogs in areas becoming drier were identified as potentially vulnerable habitats. Hodgson et al. (2009) also cite several other modelling studies focussing on Britain and examining potential impacts on butterfly and bird species.

---

<sup>15</sup> When talking about invasive alien species in the context of climate change, it is important to note that according to the CBD definition of invasive alien species as interpreted by the Standing Committee to the Bern Convention, native species naturally extending their range in response to climate change should not be considered as "alien" (Rec. No. 142 (2009)).

<sup>16</sup> Doc. T-PVS/Inf(2009)12 and T-PVS (2009) 13

<sup>17</sup> Note also Bern Convention Recommendation 91 (2002) on invasive species that threaten biological diversity on islands and evolutionary isolated ecosystems

According to Heywood & Culham (2009), the islands of Macaronesia should be considered as a particularly vulnerable region due to the potential effects of an eastwardly shift of the Azorean anticyclone that would diminish the frequency and intensity of the northwest trade winds with consequential effects on the unique Laurel forest zone. Citing an assessment of climate change impacts on the flora and vegetation of the Canary islands by Del Arco (2008), they further point out risks to high altitude ecosystems and species due to their inability to migrate.

Hodgson et al. (2009) note a very limited amount of available research on the vulnerability of island biota in the Mediterranean region, but assume a high vulnerability of their endemic species due to the geographic restrictions. Heywood & Culham also suggest that Mediterranean islands are at high risk of facing species extinctions both because of the presence of a large number of narrowly endemic and threatened plant species, many of which are confined to mountain areas making successful dispersal into new areas even more unlikely, and the significant climatic changes expected. The generally high vulnerability of Mediterranean ecosystems due to an increase in already now severe drought stress and fire risk has been observed in several studies (Berry 2008).

Based on expert judgment, it is assumed that rising temperatures and a decrease in days with precipitation could threaten the endemic Corsican gastropod *Tyrrhenaria ceratina* as well as several endemic species of Corsican mountain aquatic invertebrates (French national report to the Bern Convention).

The expected impacts of climate change in the Arctic biogeographical region, including thawing of permafrost and a major reduction in the area of tundra (EEA 2002-2008, Zockler and Lysenko 2000), are likely to affect island ecosystems in the region.

#### **4. IMPLICATIONS OF CLIMATE CHANGE FOR CONSERVATION EFFORTS ON EUROPEAN ISLANDS AND OPTIONS FOR ACTION**

Climate change adds to the existing pressures on island biodiversity and is likely to affect those species most which are often already threatened - species with specialized habitat requirements, small population numbers, limited distribution areas and low dispersal ability. Therefore, many conservation strategies that have been put in place will remain useful under conditions of climate change.

However, the increasing dynamics of environmental conditions and the possibility of unexpected effects (triggered for example by impacts of climate change on species interactions) have to be taken into account, both by stepping up monitoring efforts with a particular focus on islands and by ensuring a certain amount of flexibility to adapt strategies if needed.

In addition, targeted measures to mitigate the negative consequences of climate change should be taken. Thorough reviews of possible approaches to adaptation in biodiversity conservation have been provided by Huntley (2007) and Harley (2008). However, not all of these approaches can be applied equally well to islands.

While measures such as filling the gaps in protected area networks, maintaining and restoring ecosystem function, reducing non-climatic stresses, introducing elements of adaptive management and establishing buffering zones are relevant on islands as much as anywhere else, the insular setting effectively limits the applicability of those strategies which aim to facilitate natural adaptation by shifts in species distributions. Measures to increase the connectivity of habitats by the creation of corridors or stepping stones and by strengthening landscape permeability in general are feasible only within islands and would thus seem irrelevant to many of the smaller island territories.

Because of this limitation, and also with a view to the characteristic high level of endemism on certain islands, special consideration should be given to the question of ex situ conservation and translocation for those species which are threatened with extinction in their current habitat, and unlikely to be able to reach other suitable habitat by natural dispersal. Although both ex situ and translocation measures are very resource-intensive strategies and not always feasible in practice, and translocation also carries a significant amount of risk to biota in the target area (see also discussion in Heywood & Culham 2009 and CBD Secretariat 2009), where such options exist they may be the only way to ensure the survival of certain taxa.

The specific threats to coastal habitats from sea-level rise should be addressed by providing space for natural retreat towards higher areas wherever feasible, and by avoiding further habitat destruction for the construction of settlement and transport infrastructure.

Finally, preventive measures against the establishment of further invasive species and containment measures where invasive alien species have already been introduced should be considered a priority in island ecosystems.

## **5. RESEARCH NEEDS**

In addition to research needs already identified in previous reports (including improving the information base on the vulnerability of Bern Convention species and habitats, and strengthening monitoring schemes) and by other Expert Groups (including the identification of knowledge gaps in European island threatened biodiversity and on invasive alien species on European islands), the following specific research needs should be addressed:

- improving knowledge about island endemic species in less well researched groups,
- monitoring climate change impacts on island biota (including impacts on migratory species),
- further development of appropriate approaches to assess the vulnerability of rare and endemic species to climate change, including trait-based assessment frameworks.
- improving climate projections at a resolution which is appropriate for consideration of climate change effects on islands

## **6. CONCLUSIONS AND RECOMMENDATIONS**

Many recommendations which are relevant to the conservation of island biodiversity under climate change have already been approved by the Standing Committee and should be applied in the island context.

These include:

- Bern Convention Recommendation 135 (2008) on addressing the impacts of climate change on biodiversity, and in particular the points of guidance on taking an integrated approach to climate change response activities, addressing non-climatic threats to vulnerable species, taking early action on the protection of island-endemic amphibian and reptile species, maintaining and restoring large intact habitats as well as ecosystem structure and function, establishing networks of interconnected protected areas, increasing protected area coverage where necessary to ensure that vulnerable species groups and habitats are included, establishing buffer zones around conservation areas, avoiding development in coastal areas, considering the role of species translocation and ex situ conservation, ensuring policy integration, using adaptive management and addressing invasive species issues.
- Bern Convention Recommendation 143 (2009) on further guidance for Parties on biodiversity and climate change, and in particular the points of guidance on minimising threats to vulnerable invertebrates, including in Atlantic and Mediterranean islands, implementing appropriate protected area management to increase resilience and considering mechanisms for implementation of off-protected areas management.
- Bern Convention Recommendation 91 (2002) on invasive species that threaten biological diversity on islands and evolutionary isolated ecosystems
- The European Strategy on invasive species

When developing adaptation measures, special consideration should be given to islands of the Mediterranean and Macaronesian regions because of their high rates of endemism and expected serious changes in precipitation regimes, and within these regions particularly to those sites hosting vulnerable or threatened endemic taxa (the lists provided in annexes I and II can serve as a starting point for this). However, islands in other regions may also contain highly sensitive biota which require attention, as exemplified by the observed drastic declines in seabird populations of the North East Atlantic region.



## 7. ACKNOWLEDGEMENTS

I would like to thank Annabelle Cuttelod, Melanie Bilz (both IUCN) and Stuart Butchart (Birdlife International) for help in accessing information on island endemic species as well as related taxonomic information, as well as Monika Bertzky for information related to the analyses carried out in the EU Climate Change and Natura 2000 project.

## 8. REFERENCES

- Araújo, M. B., Thuiller, W. & Pearson, R.G. (2006). Climate warming and the decline of amphibians and reptiles in Europe. *Journal of Biogeography* 33: 1712-1728.
- Berry, P.M., Harrison, P.A., Dawson, T.P. and Walmsley, C.A. (Eds.) (2005). *Modelling Natural Resource Responses to Climate Change (MONARCH): A Local Approach*. UKCIP Technical Report, Oxford.
- Berry, P.M., Jones, A.P., Nicholls, R.J., Vos, C.C., (Eds.) 2007. *Assessment of the vulnerability of terrestrial and coastal habitats and species in Europe to climate change, Annex 2 of Planning for biodiversity in a changing climate - BRANCH project Final Report*. Natural England, UK.
- Berry, P.M. (2008). *Climate change and the vulnerability of Bern Convention species and habitats*. T-PVS/Inf (2008) 6 rev.
- Bertzky, M., Lysenko, I., Ravilious, C., Dickson, B., de Soye, Y. (2009). *Impacts of climate change and selected renewable energy infrastructures on EU biodiversity and the Natura 2000 network. Task 3a - Applying the vulnerability assessment framework: impacts of climate change on the Natura 2000 network*
- Capdevila-Arguelles, L., Zilletti, B. (2008). *A Perspective on Climate Change and Invasive Alien Species*. T-PVS/Inf(2008)05rev.
- Carnevali, L., Genovesi, P. (2009). *Toward a European information system on Invasive Alien Species in European islands*. Draft version. T-PVS/INF (2009)13.
- CBD Secretariat (2009): *Draft Findings Of The Ad Hoc Technical Expert Group on Biodiversity and Climate Change*. CBD Technical Series No. 41.
- Delanoë, O., de Montmollin, B. (1996). *Conservation of Mediterranean Island Plants. Strategy for Action*. IUCN Publication Service, Cambridge.
- EEA (2002-2008): *Europe's biodiversity – biogeographical regions and seas*. Available online at [http://www.eea.europa.eu/publications/report\\_2002\\_0524\\_154909](http://www.eea.europa.eu/publications/report_2002_0524_154909). Last accessed on 22nd May 2010.
- EEA (2007). *Europe's environment. The fourth assessment. State of the environment report No 1/2007*. Copenhagen.
- Foden, W., Mace, G., Vié, J.-C., Angulo, A., Butchart, S., DeVantier, L., Dublin, H., Gutsche, A., Stuart, S. & Turak, E. (2008). *Species susceptibility to climate change impacts*. In: Vié, J.-C., Hilton-Taylor, C. & Stuart, S.N. (eds). *The 2008 review of the IUCN red list of threatened species*. IUCN Gland, Switzerland, 12pp.
- Genovesi, P., Shine, C. (2003). *European Strategy on Invasive Alien Species. Final Version*. T-PVS (2003) 7 revised
- Harley, M. (2008). *Review of existing international and national guidance on adaptation to climate change: with a focus on biodiversity issues*. T-PVS/Inf (2008) 12 rev.
- Helcom (2007). *Climate Change in the Baltic Sea Area – HELCOM Thematic Assessment in 2007*. Balt. Sea Environ. Proc. No. 111
- Heywood, V., Culham, A. (2009). *The impacts of climate change on plant species in Europe*. T-PVS/Inf (2009) 9.

- Hodgson, N., Glen, E., Harley, M., Pooley, M., Sajwaj, T., Schiopu, R., de Soye, Y. (2009): Impacts of climate change and selected renewable energy infrastructures on EU biodiversity and the Natura 2000 network. Task 1 – Impacts of climate change on EU biodiversity: evidence and modelling results
- Hossell JE, Briggs B & Hepburn IR (2000) Climate Change and Nature Conservation: A Review of the Impact of Climate Change on UK Species and Habitat Conservation Policy. HMSO, DETR, MAFF, London.
- Huntley, B. (2007): Climatic change and the conservation of European biodiversity: Towards the development of adaptation strategies. Final version. T-PVS/Inf (2007) 3
- Huntley, B., Green, R.E., Collingham, Y.C., Willis, S.G. (2008). A Climatic Atlas of European Breeding Birds. Lynx Ediciones
- Kühn, I., K. Vohland, F. Badeck, J. Hanspach, S. Pompe, and S. Klotz.(2009). Aktuelle Ansätze zur Modellierung der Auswirkungen von Klimaveränderungen auf die biologische Vielfalt. *Natur und Landschaft* 84:1-12.
- Nordic Council of Ministers (2009): Signs of Climate Change in Nordic Nature. Tema Nord 2009:551. Copenhagen
- Orueta J.F., 2009. International efforts to conserve biological diversity in Islands. T-PVS/INF (2009)1.
- Sajwaj, T., Tucker, G., Harley, M., de Soye, Y. (2009). Impacts of climate change and selected renewable energy infrastructures on EU biodiversity and the Natura 2000 network. Task 2a – An assessment framework for climate change vulnerability: methodology and results
- Settele, J., Kudrna, O., Harpke, A., Kuhn, I., van Sway, C., Verovnik, R., Warren, M., Wiemers, M., Hanspach, J., Hickler, T., Kuhn, E., van Halder, I., Veling, K., Vliegnerhart, A., Wynhoff, I. & Schweiger, O. (2008). Climatic risk atlas of European butterflies. Pensoft, Sofia.
- Thuiller, W., Lavorel, S., Araujo, M.B., Sykes, M.T., Prentice, I.C. (2005). Climate change threats to plant diversity in Europe. *Proceedings of the National Academy of Sciences* 102, 8245-8250.
- Whittaker, R.J., Fernandez-Palacios, J.M. (2007). *Island Biogeography. Ecology, evolution and conservation*. Second edition. Oxford University Press.
- Zöckler, C. and I. Lysenko (2000). *Water Birds on the Edge: First Circumpolar Assessment of Climate Change Impact on Arctic Breeding Water Birds*. WCMC Biodiversity Series No. 11. World Conservation Monitoring Centre, Cambridge, 20pp.

**Annex 1:**

Endemic animal species in various taxonomic groups, Bern Convention species marked (data extracted from IUCN Red Lists, IUCN Species Database, BirdLife fact sheets and WCMC Species Database)

Species name	Distribution	IUCN Red List Status	Listed in Appendix II of the Bern Convention
<b>Mammals</b>			
<i>Nyctalus azoreum</i>	Azores	EN	Yes
<i>Pipistrellus maderensis</i>	Madeira, Canary islands	EN	Yes
<i>Plecotus sardus</i>	Sardinia	VU	Yes
<i>Plecotus teneriffae</i>	Canary islands	EN	Yes
<i>Acomys minous</i>	Crete	DD	--
<i>Crocidura canariensis</i>	Canary islands	EN	Yes
<i>Crocidura sicula</i>	Sicily and surrounding islands	LC	--
<i>Crocidura zimmermanni</i>	Crete	VU	--
<b>Birds</b>			
<i>Loxia scotica</i>	Great Britain	DD	Yes
<i>Serinus canaria</i>	Azores, Madeira, Canary Islands	LC	--
<i>Pterodroma madeira</i>	Madeira	EN	Yes
<i>Columba trocaz</i>	Madeira	NT	--
<i>Columba bollii</i>	Canary islands	NT	Yes
<i>Columba junoniae</i>	Canary islands	EN	Yes
<i>Apus unicolor</i>	Madeira, Canary islands	LC	Yes
<i>Saxicola dacotiae</i>	Canary islands	EN	Yes
<i>Anthus berthelotii</i>	Madeira, Canary islands	LC	Yes
<i>Fringilla teydea</i>	Canary islands	NT	Yes
<i>Sylvia melanothorax</i>	Cyprus	LC	Yes
<i>Oenanthe cypriaca</i>	Cyprus	LC	Yes
<i>Puffinus mauretanicus</i>	Balearic islands	CR	--
<i>Sitta whiteheadi</i>	Corsica	LC	Yes
<i>Regulus madeirensis</i>	Madeira	LC	Yes
<b>Reptiles</b>			
<i>Hierophis cypriensis</i> ( <i>Coluber cypriensis</i> )	Cyprus	EN	Yes
<i>Tarentola angustimentalis</i>	Canary Islands	LC	Yes
<i>Tarentola boettgeri</i>	Canary Islands	LC	Yes
<i>Tarentola delalandii</i>	Canary Islands	LC	Yes
<i>Tarentola gomerensis</i>	Canary islands	LC	Yes
<i>Algyroides fitzingeri</i>	Corsica, Sardinia and some surrounding islets	LC	Yes
<i>Archaeolacerta bedriagae</i>	Corsica, Sardinia and several smaller islands	NT	Yes
<i>Gallotia atlantica</i>	Canary Islands	LC	--
<i>Gallotia auaritae</i>	Canary Islands (possibly extinct)	CR	Yes (as part of <i>Gallotia simonyi</i> )
<i>Gallotia bravoana</i>	Canary Islands	CR	Yes (as part of <i>Gallotia</i> )

			simonyi)
<i>Gallotia caesaris</i>	Canary Islands	LC	Yes (as part of <i>Gallotia galloti</i> )
<i>Gallotia galloti</i>	Canary Islands	LC	Yes
<i>Gallotia intermedia</i>	Canary islands	CR	--
<i>Gallotia simonyi</i>	Canary Islands	CR	Yes
<i>Gallotia stehlini</i>	Canary Islands	LC	Yes
<i>Phoenicolacerta troodica</i>	Cyprus	LC	--
<i>Podarcis cretensis</i>	Crete and satellite islands	EN	Yes (as part of <i>Podarcis erhardii</i> )
<i>Podarcis filfolensis</i>	Malta, Linosa and Lampione (Italy)	LC	Yes
<i>Podarcis gaigeae</i>	Aegean Islands	VU	--
<i>Podarcis levendis</i>	Aegean Islands	VU	--
<i>Podarcis lilfordi</i>	Balearic Islands	EN	Yes
<i>Podarcis milensis</i>	Aegean islands	VU	Yes
<i>Podarcis pityusensis</i>	Balearic Islands	NT	Yes
<i>Podarcis raffonei</i>	Aeolian Islands (Italy)	CR	--
<i>Podarcis tiliguerta</i>	Corsica, Sardinia and adjacent islets	LC	Yes
<i>Podarcis waglerianus</i>	Sicily, adjacent islands	LC	Yes
<i>Teira dugesii</i>	Native to Madeira, introduced to the Azores Islands	LC	Yes
<i>Chalcides sexlineatus</i>	Canary Islands	LC	Yes
<i>Chalcides simonyi</i> ( <i>Chalcides occidentalis</i> )	Canary Islands	EN	Yes
<i>Chalcides viridanus</i>	Canary Islands	LC	Yes
<i>Macrovipera schweizeri</i> ( <i>Vipera schweizeri</i> )	Aegean Islands	EN	Yes
<i>Emys trinacris</i>	Sicily	DD	Yes (as part of <i>Emys orbicularis</i> )
<b>Amphibians</b>			
<i>Speleomantes imperialis</i>	Sardinia	NT	Yes
<i>Euproctus platycephalus</i>	Sardinia	EN	Yes
<i>Lyciasalamandra helverseni</i>	Aegean Islands	VU	--
<i>Atylodes (Hydromantes / Speleomantes) genei</i>	Sardinia	VU	Yes
<i>Alytes muletensis</i>	Balearic Islands	VU	Yes
<i>Salamandra corsica</i>	Corsica	LC	--
<i>Pseudepidalea sicula</i>	Sicily, adjacent islands	LC	Yes (as part of <i>Bufo viridis</i> )
<i>Euproctus montanus</i>	Corsica	LC	Yes
<i>Speleomantes (Hydromantes) flavus</i>	Sardinia	VU	Yes
<i>Speleomantes supramontis</i>	Sardinia	EN	Yes
<i>Pelophylax cretensis</i>	Crete	EN	--
<i>Pelophylax cerigensis</i>	Aegean Islands	CR	--
<i>Discoglossus montalentii</i>	Corsica	NT	Yes
<i>Hyla sarda</i>	Corsica, Sardinia, adjacent islands	LC	Yes
<i>Speleomantes sarrabusensis</i>	Sardinia	VU	Yes (as part of <i>Hydromantes / Speleomantes imperialis</i> )

<b>Butterflies</b>			
<i>Plebejus psyloritus</i>	Crete	LC	--
<i>Plebejus bellieri</i>	Corsica, Sardinia	LC	--
<i>Cyclus webbianus</i>	Canary Islands	LC	--
<i>Glaucopsyche paphos</i>	Cyprus	LC	--
<i>Pseudophilotes barbaggioi</i>	Sardinia	LC	--
<i>Papilio hospiton</i>	Corsica, Sardinia	LC	Yes
<i>Zerynthia cretica</i>	Crete	LC	--
<i>Thymelicus christi</i>	Canary Islands	LC	--
<i>Spialia therapne</i>	Corsica and Sardinia	LC	--
<i>Hipparchia maderensis</i>	Madeira	LC	--
<i>Hipparchia azorina</i>	Azores	LC	--
<i>Hipparchia tilosi</i>	Canary Islands	VU	--
<i>Hipparchia aristaeus</i>	Corsica, Sardinia, adjacent islands	LC	--
<i>Hipparchia gomera</i>	Canary Islands	LC	--
<i>Hipparchia neomiris</i>	Corsica, Sardinia, adjacent islands	LC	--
<i>Hipparchia sbordonii</i>	Ponza Islands, Italy	NT	--
<i>Hipparchia miguelensis</i>	Azores	LC	--
<i>Hipparchia leighebi</i>	Aeolian Islands, Italy	NT	--
<i>Hipparchia cypriensis</i>	Cyprus	LC	--
<i>Hipparchia cretica</i>	Crete	LC	--
<i>Hipparchia bacchus</i>	Canary islands	VU	--
<i>Hipparchia tamadabae</i>	Canary islands	LC	--
<i>Hipparchia christenseni</i>	Aegean islands	LC	--
<i>Hipparchia wyssii</i>	Canary islands	LC	--
<i>Coenonympha thyrus</i>	Crete	LC	--
<i>Coenonympha corinna</i>	Corsica, Sardinia and adjacent islands	LC	--
<i>Pararge xiphia</i>	Madeira	EN	--
<i>Pararge xiphioides</i>	Canary islands	LC	--
<i>Melanargia pherusa</i>	Sicily	LC	--
<i>Vanessa vulcania</i>	Madeira, Canary islands	LC	--
<i>Argynnis elisa (Fabriciana elisa)</i>	Corsica, Sardinia	LC	Yes
<i>Maniola nurag</i>	Sardinia	LC	--
<i>Maniola chia</i>	Aegean islands	LC	--
<i>Maniola cypricola</i>	Cyprus	LC	--
<i>Lasiommata paramegaera</i>	Corsica, Sardinia and adjacent islands	LC	--
<i>Aglais ichnusa</i>	Corsica, Sardinia	LC	--
<i>Gonepteryx maderensis</i>	Madeira	EN	--
<i>Gonepteryx cleobule</i>	Canary islands	VU	--
<i>Euchloe insularis</i>	Corsica, Sardinia	LC	--
<i>Euchloe grancanariensis</i>	Canary islands	LC	--
<i>Euchloe hesperidum</i>	Canary islands	LC	--
<i>Euchloe eversi</i>	Canary islands	LC	--
<i>Pieris cheiranthi</i>	Canary islands	EN	--
<i>Pieris wollastoni</i>	Madeira	CR	--
<b>Dragonflies</b>			
<i>Boyeria cretensis</i>	Crete	EN	--
<i>Sympetrum nigrifemur</i>	Canary Islands, Madeira	LC	--
<i>Coenagrion intermedium</i>	Crete	VU	--
<i>Ischnura genei</i>	Corsica, Sardinia, Sicily, Malta and adjacent islands	LC	--

Saproxyllic Beetles			
<i>Trichoferus bergeri</i>	Crete	EN	--
<i>Axinopalpis barbarae</i>	Cyprus	DD	--
<i>Delagrangaeus schurmanni</i>	Canary Islands	VU	--
<i>Crotchiella brachyptera</i>	Azores	EN	--
<i>Stenopterus creticus</i>	Crete	EN	--
<i>Blabinotus spinicollis</i>	Canary Islands, Madeira	NT	--
<i>Glaphyra bassettii</i>	Cyprus	CR	--
<i>Pseudosphegesthes bergeri</i>	Crete	EN	--
<i>Anaglyptus praecellens</i>	Crete	EN	--
<i>Purpuricenus nicocles</i>	Cyprus	NT	--
<i>Isotomus jarmilae</i>	Crete	EN	--
<i>Clytus clavicornis</i>	Sicily	VU	--
<i>Pediacus tabellatus</i>	Canary islands	DD	--
<i>Scobicia barbifrons</i>	Canary Islands	LC	--
<i>Scobicia ficicola</i>	Canary Islands	LC	--
<i>Stephanopachys brunneus</i>	Canary Islands	NT	--
<i>Procraterus cretensis</i>	Crete	DD	--
<i>Stenagostus sardiniensis</i>	Sardinia, possibly Corsica	DD	--
<i>Ampedus minos</i>	Crete	DD	--
<i>Ampedus assingi</i>	Cyprus	EN	--
<i>Ampedus corsicus</i>	Corsica	NT	--
<i>Alestrus dolosus</i>	Azores	DD	--
<i>Haterumelater schembrii</i>	Malta	DD	--
<i>Gnorimus decempunctatus</i>	Sicily	VU	--
<i>Protaetia sardea</i>	Sardinia and Corsica	DD	--
<i>Osmoderma cristinae</i>	Sicily	EN	Yes (as part of <i>Osmoderma eremita</i> )
<i>Leipaspis pinicola</i>	Canary Islands	VU	--
<i>Leipaspis lauricola</i>	Canary Islands	VU	--
<i>Propomacrus cypriacus</i>	Cyprus	CR	--
<i>Dorcus musimon</i>	Corsica, Sardinia	LC	--
<i>Dorcus alexisi</i>	Cyprus	EN	--

**Annex 2:**

Plant species protected under the Bern Convention which are endemic to a single island or archipelago in Europe according to the Euro+Med Plantbase<sup>18</sup>

<b>Species name</b>	<b>Distribution</b>
<i>Primula egaliksensis</i>	Iceland
<i>Bromus interruptus</i>	Great Britain
<i>Gentianella anglica</i>	Great Britain
<i>Anthyllis hystrix</i>	Balearic islands
<i>Centaurea balearica</i>	Balearic islands
<i>Daphne rodriguezii</i>	Balearic islands
<i>Diploxix ibicensis</i>	Balearic islands
<i>Euphorbia margalidiana</i>	Balearic islands
<i>Genista dorycnifolia</i>	Balearic islands
<i>Lysimachia minoricensis</i>	Balearic islands
<i>Paeonia cambessedesii</i>	Balearic islands
<i>Ranunculus weyleri</i>	Balearic islands
<i>Viola jaubertiana</i>	Balearic islands
<i>Aconitum corsicum</i>	Corsica
<i>Astragalus maritimus</i>	Sardinia
<i>Centaurea horrida</i>	Sardinia
<i>Lamyropsis microcephala</i>	Sardinia
<i>Ribes sardoum</i>	Sardinia
<i>Abies nebrodensis</i>	Sicily
<i>Brassica macrocarpa</i>	Sicily
<i>Bupleurum dianthifolium</i>	Sicily
<i>Cytisus aeolicus</i>	Sicily
<i>Petagnaena saniculifolia</i>	Sicily
<i>Cremnophyton lanfrancoi</i>	Malta
<i>Helichrysum melitense</i>	Malta
<i>Palaeocyanus crassifolius</i>	Malta
<i>Asyneuma giganteum</i>	Aegean islands
<i>Consolida samia</i>	Aegean islands
<i>Androcymbium rechingeri</i>	Crete
<i>Anthemis glaberrima</i>	Crete
<i>Arum purpureospathum</i>	Crete
<i>Bupleurum kakiskalae</i>	Crete
<i>Carlina diae</i>	Crete
<i>Cephalanthera cucullata</i>	Crete
<i>Colchicum cousturieri</i>	Crete
<i>Convolvulus argyrothamnos</i>	Crete
<i>Hypericum aciferum</i>	Crete
<i>Origanum dictamnus</i>	Crete
<i>Wagenitzia lancifolia</i>	Crete
<i>Zelkova abelicea</i>	Crete
<i>Alyssum akamasicum</i>	Cyprus
<i>Arabis kennedyae</i>	Cyprus
<i>Brassica hilarionis</i>	Cyprus
<i>Centaurea akamantis</i>	Cyprus
<i>Chionodoxa lochiai</i>	Cyprus

<sup>18</sup> Endemic species of the Macaronesian region were purposely omitted from this table as they are already listed separately in Appendix I of the Convention.

<i>Crocus cyprius</i>	Cyprus
<i>Crocus hartmannianus</i>	Cyprus
<i>Delphinium caseyi</i>	Cyprus
<i>Onosma troodi</i>	Cyprus
<i>Ophrys kotschyi</i>	Cyprus
<i>Origanum cordifolium</i>	Cyprus
<i>Phlomis brevibracteata</i>	Cyprus
<i>Phlomis cypria</i>	Cyprus
<i>Ranunculus kykkoensis</i>	Cyprus
<i>Salvia veneris</i>	Cyprus
<i>Scilla morrisii</i>	Cyprus
<i>Sideritis cypria</i>	Cyprus
<i>Tulipa cypria</i>	Cyprus



**Annex 3:**

Species protected under Appendix I or II of the Bern Convention which are assessed by Sajwaj et al. (2009) as being at least highly vulnerable to climate change.

<i>Accipiter brevipes</i>	<i>Nycticorax nycticorax</i>
<i>Acrocephalus melanopogon</i>	<i>Oceanodroma leucorhoa</i>
<i>Acrocephalus paludicola</i>	<i>Oenanthe leucura</i>
<i>Aegypius monachus</i>	<i>Oenanthe pleschanka</i>
<i>Anser erythropus</i>	<i>Otis tarda</i>
<i>Anthus campestris</i>	<i>Oxyura leucocephala</i>
<i>Apus caffer</i>	<i>Pandion haliaetus</i>
<i>Aquila adalberti</i>	<i>Phalacrocorax pygmeus</i>
<i>Aquila chrysaetos</i>	<i>Phengaris nausithous (Maculinea nausithous)</i>
<i>Aquila clanga</i>	<i>Phoenicopterus ruber</i>
<i>Aquila heliaca</i>	<i>Picoides tridactylus</i>
<i>Aquila pomarina</i>	<i>Picus canus</i>
<i>Ardea purpurea</i>	<i>Platalea leucorodia</i>
<i>Ardeola ralloides</i>	<i>Plegadis falcinellus</i>
<i>Asio flammeus</i>	<i>Podiceps auritus</i>
<i>Botaurus stellaris</i>	<i>Porphyrio porphyrio</i>
<i>Botrychium simplex</i>	<i>Porzana parva</i>
<i>Branta leucopsis</i>	<i>Porzana pusilla</i>
<i>Bubo bubo</i>	<i>Pterocles alchata</i>
<i>Burhinus oediconemus</i>	<i>Pterocles orientalis</i>
<i>Buteo rufinus</i>	<i>Puffinus yelkouan</i>
<i>Calonectris diomedea (Procellaria diomedea)</i>	<i>Pulsatilla patens</i>
<i>Charadrius morinellus (Eudromias morinellus)</i>	<i>Pyrhocorax pyrrhocorax</i>
	<i>Rana latastei</i>
	<i>Recurvirostra avosetta</i>
	<i>Rumex rupestris</i>
<i>Chersophilus duponti</i>	<i>Salamandrina terdigitata</i>
<i>Chioglossa lusitannica</i>	<i>Sitta krueperi</i>
<i>Chlidonias hybridus</i>	<i>Sterna albifrons</i>
<i>Chlidonias niger</i>	<i>Sterna caspia (Hydroprogne caspia)</i>
<i>Ciconia nigra</i>	<i>Sterna dougallii</i>
<i>Circaetus gallicus</i>	<i>Sterna paradisaea</i>
<i>Circus cyaneus</i>	<i>Sterna sandvicensis</i>
<i>Circus macrourus</i>	<i>Strix nebulosa</i>
<i>Circus pygargus</i>	<i>Strix uralensis</i>
<i>Coracias garrulus</i>	<i>Surnia ulula</i>
<i>Crex crex</i>	<i>Sylvia rueppelli</i>
<i>Cygnus bewickii (Cygnus columbianus bewickii)</i>	<i>Sylvia sarda</i>
<i>Cygnus cygnus</i>	<i>Sylvia undata</i>
<i>Dendrocopos leucotos</i>	<i>Tadorna ferruginea</i>
<i>Dendrocopos medius</i>	<i>Tetrax tetrax</i>
<i>Discoglossus galganoi</i>	<i>Triturus montandoni</i>
<i>Egretta alba (Ardea alba) (Casmerodius albus)</i>	<i>Vipera ursinii</i>
<i>Egretta garzetta</i>	<i>Xenus cinereus (Tringa cinerea)</i>
<i>Elanus caeruleus</i>	
<i>Emberiza cineracea</i>	
<i>Falco biarmicus</i>	
<i>Falco cherrug</i>	
<i>Falco columbarius</i>	
<i>Falco eleonora</i>	
<i>Falco rusticolus</i>	

*Falco vespertinus*  
*Ficedula albicollis*  
*Ficedula semitorquata*  
*Fulica cristata*  
*Galerida theklae*  
*Gallinago media*  
*Gelochelidon nilotica* (*Sterna nilotica*)  
*Glareola pratincola*  
*Grus grus*  
*Gypaetus barbatus*  
*Gyps fulvus*  
*Haliaeetus albicilla*  
*Hieraaetus fasciatus*  
*Hieraaetus pennatus*  
*Himantopus himantopus*  
*Hydrobates pelagicus*  
*Lacerta monticola* (*Archaeolacerta monticola*)  
*Lacerta schreiberi*  
*Lanius nubicus*  
*Larus audouinii*  
*Larus melanocephalus*  
*Loxia scotica*  
*Luscinia svecica*  
*Lycaena dispar*  
*Marmaronetta angustirostris*  
*Melanargia arge*  
*Mergus albellus* (*Mergellus albellus*)  
*Milvus milvus*  
*Neophron percnopterus*  
*Nyctea scandiaca*